Author Index

Ashby, J.M.: See Li, Y., et al., 26
Atkin, J.D.: See Bender, E., et al., 74

Baba, M., Kobayashi, T., Oka, Y., Yagyu, T., Tamaki, Y., Takeda, T., Monden, T., Shimano, T., Tsuji, Y., Murakami, H.: Preparation of a human monoclonal antibody derived from cervical lymph nodes of a patient with anaplastic carcinoma of the thyroid, 181
Barker, M.D.: See Bender, E., et al., 74
Bebbington, C.R.: See Bender, E., et al., 74
Bell, D.A.: See Denomme, G.A., et al., 98
Bende, R.J.: See Jochems, G.J., et al., 124
Bender, E., Woof, J.M., Atkin, J.D., Barker, M.D., Bebbington, C.R., Burton, D.R.: Recombinant human antibodies: linkage of an Fab fragment from a combinational library to an Fc fragment for expression in mammalian cell culture, 74
Bitoh, S., Lang, G.M., Sehon, A.H.: Suppression of human anti-mouse idiotypic antibody responses in hu-PBL-SCID mice, 144
Bitoh, S., Lang, G.M., Kierek-Jaszczuk, D., Fujimoto, S., Sehon, A.H.: Specific immunosuppression of human anti-murine antibody (HAMA) responses in hu-PBL-SCID mice, 134
Boerner, P.: See Brams, P., et al., 47, 57
Borrebaeck, C.A.K.: See Ifversen, P., et al., 115
Brams, P., Royston, I., Boerner, P.: In vitro priming of human lymphocytes. I. IL-2 and IL-4 requirements, 47
Brams, P., Royston, I., Boerner, P.: In vitro priming of human lymphocytes. II. Induction of antigen-specific IgG responses by repeated antigen stimulation, 57
Burton, D.R.: See Bender, E., et al., 74

Cairns, E.: See Denomme, G.A., et al., 98
Cartner, A.M., Conry, R.M., Safavy, A., Khazaeli, M.B., Sumerel, L.A., LoBuglio, A.F.: An animal model to predict the immunogenicity of murine V-regions in humans, 174
Conry, R.M.: See Cartner, A.M., et al., 174

Daifuku, R., Panacek, E.A., Haenftling, K., Swenson, W.K., Prescott, A.W., Johnson, J.L.: Pilot study of anti-lipopolysaccharide human monoclonal antibody MAB-T88 in patients with gram-negative sepsis, 36
Dang, H.: See Lazaridis, K., et al., 104
de Groot, C.: See Mevissen, M.C.L.M., et al., 66
Denomme, G.A., Mahmoudi, M., Edwards, J.Y., Massicotte, H., Cairns, E., Bell, D.A.: Immunoglobulin V region heavy and light chain gene sequences of the lymphoblastoid cell line GM 4672, 98
Ditzel, H., Erb, K., Leslie, G., Jensenius, J.C.: Preparation of antigen-binding monomeric and half-monomeric fragments from human monoclonal IgM antibodies against colorectal cancer-associated antigens, 86

Edwards, J.Y.: See Denomme, G.A., et al., 98
Erb, K.: See Ditzel, H., et al., 86
Eremin, O.: See Li, Y., et al., 26

Feng, O.: See Liu, H., et al., 2
Ferraro, A.S., Newkirk, M.M.: In vitro stimulation of human peripheral blood B cells from normal individuals by activated T cells increases the efficiency of hybridoma generation, 80
Fujimoto, S.: See Bitoh, S., et al., 134

Glassy, M.C.: Production methods for generating human monoclonal antibodies, 154
Gonzalez, M.F.: See Sanchez, B., et al., 198
Goodall, M.: See Lund, J., et al., 20
Goormachtig, E.: See Mevissen, M.L.C.M., et al., 66
Grizzle, W.E.: See Meredith, R.F., et al., 190

Haenftling, K.: See Daifuku, R., et al., 36
Hagiwara, H., Yuasa, H.: Generation of somatic cell hybrids capable of proliferating and secreting human monoclonal antibody without any growth factor supplements, 15
Hashizume, S.: See Kato, M., et al., 9
Hashizume, S.: See Yasumoto, K., et al., 186
Hindley, S.: See Lund, J., et al., 20
Huang, C.: See Zhu, Y., et al., 31

Ifversen, P., Xiu-Mei, Z., Ohlin, M., Zeuthen, J., Borrebaeck, C.A.K.: Effect of cell derived growth factors and cytokines on the clonal outgrowth of EBV infected-B cells and established lymphoblastoid cell lines (LCL), 115

Jefferis, R.: See Lund, J., et al., 20
Jensenius, J.C.: See Ditzel, H., et al., 86
Jin, B.: See Zhu, Y., et al., 31
Jochems, G.J., Bende, R.J., Klein, M.R., Zeijlemaker, W.P., van Lier, R.A.W.: Expression of the II-6 gene induces differentiation of a human monoclonal EBV transformed cell line, 124
Johnson, J.L.: See Daifuku, R., et al., 36

Kamei, M.: See Yasumoto, K., et al., 186
Kato, M.: See Yasumoto, K., et al., 186
Kato, M., Mochizuki, K., Hashizume, S., Tachibana, H., Shirahata, S., Murakami, H.: Activity enhancement of a lung cancer-associated human monoclonal antibody HB4C5 by *N*-deglycosylation, 9
Khazaeli, M.B.: See Cartner, A.M., et al., 174
Khazaeli, M.B.: See Meredith, R.F., et al., 190
Kierek-Jaszczuk, D.: See Bitoh, S., et al., 134
Klein, M.R.: See Jochems, G.J., et al., 124
Kobayashi, Tetsuro: See Baba, M., et al., 181
Kwekkeboom, J.: See Mevissen, M.L.C.M., et al., 66

Lang, Glen M.: See Bitoh, S., et al., 144
Lang, Glen M.: See Bitoh, S., et al., 134
Lazaridis, K., Dang, H., Talal, N.: Human K chain expression in a phage vector. methods of isolating amplified cDNA affects cloning efficiency, 104
Leslie, G.: See Ditzel, H., et al., 86
Li, Y.: See Liu, H., et al., 2
Li, Y., Ashby, J.M., Eremin, O.: In vitro primary immunization of B lymphocytes for producing human monoclonal antibodies against tumor-associated antigens, 26–30
Lindhout, E.: See Mevissen, M.L.C.M., et al., 66
Liu, H., Xu, Z.-L., Wang, Y., Yang, L., Feng, O., Li, Y., Wang, Y.-M., Zhang, G.-G.: Production of anti-tumor human monoclonal antibodies using different approaches, 2–8
Liu, T.: See Meredith, R.F., et al., 190
Liu, X.: See Zhu, Y., et al., 31
LoBuglio, A.F.: See Cartner, A.M., et al., 174
LoBuglio, A.F.: See Meredith, R.F., et al., 174
Lund, J., Takahashi, N., Hindley, S., Tyler, R., Goodall, M., Jefferis, R.: Glycosylation of human IgG subclass and mouse IgG2b heavy chains secreted by mouse J558L transfectoma cell lines as chimeric antibodies, 20

Mahmoudi, M.: See Denomme, G.A., et al., 98
Massicotte, H.: See Denomme, G.A., et al., 98
Melero, J.: See Sanchez, B., et al., 198
Meredith, R.F., Khazaeli, M.B., Grizzle, W.E., Orr, R.A., Plott, G., Urist, M.M., Liu, T., Russell, C.D., Wheeler, R.H., Schlom, J., LoBuglio, A.F.: Direct localization comparison of murine and chimeric B72.3 antibodies in patients with colon cancer, 190
Mevissen, M.L.C.M., Kwekkeboom, J., Goormachtig, E., Lindhout, E., de Groot, C.: Improvement of EBV transformation and cloning efficiency of human B cells using culture supernatants from lymphoblastoid cell lines, 66
Mochizuki, K.: See Kato, M., et al., 9
Monden, T.: See Baba, M., et al., 181
Mori, T.: See Baba, M., et al., 181
Murakami, H.: See Kato, M., et al., 9
Murakami, H.: See Tachibana, H., et al., 42
Murakami, H.: See Yasumoto, K., et al., 186
Murakami, H.: See Baba, M., et al., 181

Newkirk, M.M.: See Ferraro, A.S., 80
Newkirk, M.M.: See Rioux, J.D., et al., 107
Nomoto, K.: See Yasumoto, K., et al., 186

Ohlin, M.: See Ifversen, P., et al., 115
Oka, Y.: See Baba, M., et al., 181
Orr, R.A.: See Meredith, R.F., et al., 190

Panacek, E.A.: See Daifuku, R., et al., 36
Plott, G.: See Meredith, R.F., et al., 190
Prescott, A.W.: See Daifuku, R., et al., 36

Rauch, J.: See Rioux, J.D., et al., 107
Rioux, J D., Rauch, J., Zdarsky, E., Newkirk, M.M.: Molecular characterization of the GM 4672 human lymphoblastoid cell line and analysis of its

use as a fusion partner in the generation of human-human hybridoma autoantibodies, 107
Robledo, M.M.: See Sanchez, B., et al., 198
Royston, I.: See Brams, P., et al., 47, 57
Russell, C.D.: See Meredith, R.F., et al., 190

Safavy, A.: See Cartner, A.M., et al., 174
Sanchez, B., Melero, J., Robledo, M.M., Tarrago, D., Yelamos, J., Gonzalez, M.F.: Application of cellular Elisa (CELISA) to the detection of human monoclonal autoantibodies, 198
Schlom, J.: See Meredith, R.F., et al., 190
Sehon, A.H.: See Bitoh, S., et al., 134
Sehon, A.H.: See Bitoh, S., et al., 144
Setoguchi, Y.: See Yasumoto, K., et al., 186
Shimano, T.: See Baba, M., et al., 181
Shirahata, S.: See Kato, M., et al., 9
Shirahata, S.: See Tachibana, H., et al., 42
Steenbakkers, P.G.A., Van Wezenbeck, P.M.G.F., van Zanten, J., The, T.H.: Efficient generation of human anti-cytomegalovirus IgG monoclonal antibodies from preselected antigen-specific B cells, 166
Sumerel, L.A.: See Cartner, A.M., et al., 174
Sun, C.: See Zhu, Y., et al., 31
Swenson, W.K.: See Daifuku, R., et al., 36

Tachibana, H.: See Kato, M., et al., 9
Tachibana, H., Shirahata, S., Murakami, H.: Human bifunctional antibody generated by heterologous association of heavy and light chains, 42
Takahashi, N.: See Lund, J., et al., 20
Takeda, T.: See Baba, M., et al., 181
Talal, N.: See Lazaridis, K., et al., 104
Tamaki, Y.: See Baba, M., et al., 181
Tarrago, D.: See Sanchez, B., et al., 198
The, T.H.: See Steenbakkers, P.G.A., et al., 166
Tsuji, Y.: See Baba, M., et al., 181
Tyler, R.: See Lund, J., et al., 20

Urist, M.M.: See Meredith, R.F., et al., 190

van Lier, R.A.W.: See Jochems, G.J., et al., 124
Van Wezenbeck, P.M.G.F.: See Steenbakkers, P.G.A., et al., 166
van Zanten, J.: See Steenbakkers, et al., 166

Wang, Y.: See Liu, H., et al., 2
Wang, Y.-M.: See Liu, H., et al., 2
Wheeler, R.H.: See Meredith, R.F., et al., 190
Woof, J.M.: See Bender, E., et al., 74

Xiu-Mei, Z.: See Ifversen, P., et al., 115
Xu, Z.-L.: See Liu, H., et al., 2

Yagyu, Toshio: See Baba, M., et al., 181
Yang, L.: See Liu, H., et al., 2
Yasumoto, K., Setoguchi, Y., Kamei, M., Kato, M., Nomoto, K., Murakami, H., Hashizume, S.: Cancer-specific binding of a mouse MoAb vs. Candida krusei cytochrome c: an antigen recognized by a cancer-associated human MoAb HB4C5, 186
Yelamos, J.: See Sanchez, B., et al., 198
Yuasa, and Yuasa, H.: See Hagiwara, H., 15

Zdarsky, E.: See Rioux, J.D., et al., 107
Zeijlemaker, W.P.: See Jochems, G.J., et al., 124
Zeuthen, J.: See Ifversen, P., et al., 115
Zhang, G.-G.: See Liu, H., et al., 2
Zhu, Y., Jin, B., Sun, C., Huang, C., Liu, X.: The effects of hybridoma growth factor in conditioned media upon the growth, cloning, and antibody production of heterohybridoma cell lines, 31

Subject Index

Activation, 9
Anaplastic carcinoma, 181
Antibody, 66, 174, 190
Antibody secretion, 31
Anti-CD3 activated T cells, 80
Anti-idiotype, 144
Anti-tumor monoclonal antibody, 2
Antigen-binding, 86
Antigen binding site, 9
Antigen-specific B cells, 166
Autoantibody, 198
Autocytotoxicity, 198

Bifunctional antibody, 42
Breast cancer, 26

Cancer-associated antigen, 186
Candida krusei cytochrome c, 186
Cell hybridization, 107
Chimera, 190
Chimeric IgG, 20
CHO cells, 74
Cloning, 31
CMV B cell clone, 166
Colorectal cancer, 86
Conditioned media, 31
Cytokines, 66

DNA isolation, 104

EBV-infection, 115
EBV hybridoma, 2
ELISA, 198
Epstein-Barr virus transformation, 66

Fragment, 86
Fusion partner, 107

Glycosylation control, 20
Gram-negative, 36

Half-monomeric, 86
HAMA, 134, 144
Heterogeneity, 9
Heterohybridoma, 31
Heterologous association, 42
Human autoantibodies, 107
Human antibody, 74
Human B cells, 66
Human × human × human × hybridoma, 15
Human-human hybridoma, 2, 181
Human hybridoma, 107
Human lymphocytes, 47
Human monoclonal antibodies, 80
Human monoclonal antibody, 2, 15, 26, 42, 86, 166, 198
Human monoclonal antibody HB4C5, 186
Human-mouse hybridoma, 2
Human spleen cells, 57
hu-PBL-SCID mice, 134
Hybridoma growth factor, 31

IgG responses, 57
IgM, 36
IL-2, 47
IL-4, 47
Immune response, 174
Immunoglobulin repertoire cloning, 74
Immunoglobulin sequence, 107
Immunohistochemistry, 181
Immunostaining, 186
Immunosuppression, 144
In vitro immunization, 26, 47
In vitro priming, 57

Light chain, 42
Light chain, 9
Lipopolysaccharide, 36
Local Immunotherapy, 181
Localization, 190
Lung cancer, 186
Lymphokine requirements, 47

Mini-electrofusion, 166
Monoclonal antibodies, 107, 174
Monoclonal antibody, 36, 181
Monoclonal antibody HB4C5, 9
Monomeric, 86
MP6, 115
mPEG conjugate, 134

N-glycosylation, 9
No keywords, 124

Polymerase chain reaction, 104
Polyreactivity, 198

Quantitation, 190

Rabbit, 174
Radiolabel, 190
Recombinant antibody, 104
Restimulation, 57

SCID mice, 144
Selection, 166
Sepsis, 36
Serum-free medium, 15

Tetanus toxoid, 74
Thioredoxin, 115
Thyroid cancer, 181
Tolerance, 134, 144
Trioma, 15
Tumor-associated antigen, 26
Tumor cell antigen, 15